

# *Arenaria bertolonii*

## Habitus and growth type

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **4**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.63**

Disturbance frequency (herb layer): **1.45**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.02**

Grazing pressure: **0.17**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*KC \(ASP\) \*Asplenietea trichomanis\*\*\*](#),  
[\*\*KE \(THL\) \*Thlaspietea rotundifolii\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**