

# *Arenaria obtusiflora*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **12.99**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **1**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Disturbance indicator values

Disturbance frequency: **2.06**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.8**

Disturbance severity (herb layer): **0.8**

Mowing frequency: **0.11**

Grazing pressure: **0.42**

Soil disturbance: **0.3**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**