# *Arisarum proboscideum*

### Habitus and growth type

Plant height [m]: **0.13**
Life span: **Perennial**
Life form: **Geophyte**

### Flower

Flowering period: **January-May, September-December**

### Fruit, seed and dispersal

Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **6**
Temperature indicator value: **7**
Moisture indicator value: **4**
Reaction indicator value: **4**
Nutrient indicator value: **5**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.05**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.15**
Mowing frequency: **0**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Grazing pressure: **0.18**
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **T1G Alnus cordata forest**
Constant species of EUNIS habitats: **T1G Alnus cordata forest**

### Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*