

Armeria pungens

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.66**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **9.5**

Moisture indicator value: **4.5**

Nutrient indicator value: **4**

Salinity indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **1.63**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0**

Grazing pressure: **0.19**

Soil disturbance: **0.53**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JB \(CRI\) *Crithmo-Staticetea*](#), [JD \(AMM\) *Ammophiletea arundinaceae*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Coastal cliff**