

# *Armeria rumelica*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **2.59**

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Moisture indicator value: **3.6**

Disturbance indicator values

Disturbance frequency: **1.68**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.56**

Grazing pressure: **0.33**

Soil disturbance: **0.24**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FG \(SAC\) \*Stipo giganteae-Agrostietea castellanae\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**