Armeria ruscinonensis

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **5**
Nutrient indicator value: **4**

Disturbance indicator values

Disturbance frequency: **0.83**
Disturbance frequency (herb layer): **0.86**
Disturbance severity: **0.37**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.26**
Grazing pressure: **0.25**
Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***

Broad habitat

Occurrence in broad habitats: **Coastal cliff**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz