

# *Artemisia pontica*

## Habitus and growth type

Plant height [m]: **0.49**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **July-October**

## Fruit, seed and dispersal

Seed mass [mg]: **1.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.8**

Moisture indicator value: **3.3**

Reaction indicator value: **7.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.92**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **0.61**

Grazing pressure: **0.31**

Soil disturbance: **0.16**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Synanthropic**

## **Distribution**

Continentality: **8**

Continentality amplitude: **5**