Asphodeline taurica

Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

Fruit, seed and dispersal

Seed mass [mg]: **5.48**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.8**
Moisture indicator value: **4.5**

Disturbance indicator values

Disturbance frequency: **2.11**
Disturbance frequency (herb layer): **2.11**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.39**
Mowing frequency: **0.24**
Grazing pressure: **0.36**
Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KH (DRY) *Drypidetea spinosae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T38 Mediterranean montane Cedrus forest**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz