

# *Astragalus penduliflorus*

## Habitus and growth type

Plant height [m]: **0.5**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **6.91**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **2.3**

Moisture indicator value: **3.7**

Reaction indicator value: **7**

Nutrient indicator value: **1.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.8**

Disturbance frequency (herb layer): **1.71**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.16**

Grazing pressure: **0.26**

Soil disturbance: **0.11**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

## **Distribution**

Continentality: **7**

Continentality amplitude: **5**