# *Astragalus subuliformis*

### Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

### Taxon origin

Origin in Europe: **native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Mesic**

#### Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.6**
Moisture indicator value: **4.8**

#### Disturbance indicator values

Disturbance frequency: **2.02**
Disturbance frequency (herb layer): **2.02**
Disturbance severity: **0.45**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.48**
Grazing pressure: **0.43**
Soil disturbance: **0.2**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops**
Constant species of EUNIS habitats: **R15 Continental dry rocky steppic grassland**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

**and dwarf scrub on chalk outcrops**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*