

Astrantia minor

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.14**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **3.3**

Moisture indicator value: **4**

Reaction indicator value: **3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.22**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.01**

Grazing pressure: **0.21**

Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [AC \(LOI\) *Loiseleurio procumbentis*-*Vaccinietea*](#), [BA \(PIC\) *Vaccinio-Piceetea*](#)

Broad habitat

Occurrence in broad habitats: **Spring, Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**