

Atriplex aucheri

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **1.53**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.19**

Grazing pressure: **0.37**

Soil disturbance: **0.35**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [EC \(KAL\) *Kalidietea foliati*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation](#), [R64 Semi-desert salt pan](#), [S94 Semi-desert riparian scrub](#)

Constant species of EUNIS habitats: [R64 Semi-desert salt pan](#), [S94 Semi-desert riparian scrub](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Saline vegetation, Scrub**