

Atriplex glabriuscula

Habitus and growth type

Plant height [m]: **0.43**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **16.72**

Fruit, seed and dispersal

Seed mass [mg]: **2.69**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **5**

Moisture indicator value: **6.3**

Reaction indicator value: **7**

Nutrient indicator value: **8**

Salinity indicator value: **4.3**

Disturbance indicator values

Disturbance frequency: **2**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.58**

Mowing frequency: **0.06**

Grazing pressure: **0.16**

Soil disturbance: **0.62**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JC \(CAK\) *Cakiletea maritima*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N11 Atlantic, Baltic and Arctic sand beach](#),
[N21 Atlantic, Baltic and Arctic coastal shingle beach](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle**

Distribution

Continentality: **3**

Continentality amplitude: **3**