

Avena fatua

Habitus and growth type

Plant height [m]: **0.73**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **21.78**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **17.11**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Moisture indicator value: **4.5**

Reaction indicator value: **6.9**

Nutrient indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.75**

Disturbance frequency (herb layer): **2.04**

Disturbance severity: **0.85**

Disturbance severity (herb layer): **0.85**

Mowing frequency: **0.44**

Grazing pressure: **0.14**

Soil disturbance: **0.79**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality: **8**

Continentality amplitude: **2**