

Ballota pseudodictamnus

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **2.25**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **9.5**

Moisture indicator value: **3**

Reaction indicator value: **7.5**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.83**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.18**

Grazing pressure: **0.33**

Soil disturbance: **0.31**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) *Ononido-Rosmarinetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 *Phoenix theophrasti* vegetation](#)

Constant species of EUNIS habitats: [T25 *Phoenix theophrasti* vegetation](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**