Barbarea bracteosa

Habitus and growth type

Plant height [m]: **0.39**
Life span: **Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **5.3**
Moisture indicator value: **6**
Reaction indicator value: **5.5**
Nutrient indicator value: **6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.12**
Disturbance frequency (herb layer): **1.64**
Disturbance severity: **0.56**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.65**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.4**
Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz