

# ***Bellium bellidioides***

## **Habitus and growth type**

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **March-August**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7**

Moisture indicator value: **4.5**

Reaction indicator value: **2.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.48**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.09**

Grazing pressure: **0.26**

Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FQ \(GEN\) \*Carici-Genistetetea lobelii\*](#),  
[KB \(POD\) \*Polypodietea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Constant species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#), [S74 Central Mediterranean mountain hedgehog-heath](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Wetland, Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**