Bellium minutum

Habitus and growth type

Plant height [m]: **0.03**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **10.5**
Moisture indicator value: **1.5**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **3**

Disturbance indicator values

Disturbance frequency: **0.79**
Disturbance frequency (herb layer): **0.86**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.15**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.19**
Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JA (SAG) *Saginetea maritimae***

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Saline vegetation, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz