

Berberis aetnensis

Habitus and growth type

Plant height [m]: **0.63**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **11.48**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **4**

Moisture indicator value: **2**

Reaction indicator value: **4**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.74**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.01**

Grazing pressure: **0.23**

Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FN \(RUM\) Rumici-Astragaletea siculi](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Constant species of EUNIS habitats: [R1J Cyrno-Sardean oromediterranean siliceous dry grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),
Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**