

# *Bifora testiculata*

## Habitus and growth type

Plant height [m]: **0.3**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **6.1**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **3.7**

Reaction indicator value: **7.7**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **2.25**

Disturbance frequency (herb layer): **2.25**

Disturbance severity: **0.92**

Disturbance severity (herb layer): **0.92**

Mowing frequency: **0.31**

Grazing pressure: **0.11**

Soil disturbance: **0.81**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**