Brassica barrelieri

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.56**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **8.1**
Moisture indicator value: **2.8**
Reaction indicator value: **4.8**
Nutrient indicator value: **3.6**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.35**
Disturbance frequency (herb layer): **1.74**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.22**
Grazing pressure: **0.25**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz