

# ***Brassica montana***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Ellenberg-type indicator values

Temperature indicator value: **9**

Disturbance indicator values

Disturbance frequency: **1.14**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.63**

Mowing frequency: **0.27**

Grazing pressure: **0.29**

Soil disturbance: **0.35**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [\*\*JB \(CRI\) \*Crithmo-Staticetea\*, KE \(THL\) \*Thlaspietea rotundifolii\*\*\*](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**