

# *Brassica napus*

## Habitus and growth type

Plant height [m]: **0.94**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **24.81**

## Flower

Flowering period: **February-October**

## Fruit, seed and dispersal

Seed mass [mg]: **3.85**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Archeophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **6.4**

Moisture indicator value: **4.9**

Reaction indicator value: **7.2**

Nutrient indicator value: **7.8**

Salinity indicator value: **0.3**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [V11 Intensive unmixed crops](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**