

# ***Brassica tournefortii***

## **Habitus and growth type**

Plant height [m]: **0.4**

Life span: **Annual**

Life form: **Therophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **10.61**

## **Flower**

Flowering period: **January-May**

## **Fruit, seed and dispersal**

Seed mass [mg]: **1.4**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **8.7**

Temperature indicator value: **10.7**

Moisture indicator value: **3.3**

Reaction indicator value: **8**

Nutrient indicator value: **5.7**

Salinity indicator value: **1**

### **Disturbance indicator values**

Disturbance frequency: **1.11**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.13**

Grazing pressure: **0.22**

Soil disturbance: **0.42**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#), [QC \(CHE\) \*Chenopodietea\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Synanthropic**