Bromopsis cappadocica

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **5.5**
Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **0.76**
Disturbance frequency (herb layer): **1.66**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.25**
Grazing pressure: **0.3**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S75 Eastern Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **S75 Eastern Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Oromediterranean scrub, Forest, Sparsely vegetated (incl. rock and scree)

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz