Bromus intermedius

Habitus and growth type

Plant height [m]: **0.41**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.36**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **7.5**
Moisture indicator value: **3**
Reaction indicator value: **7.7**
Nutrient indicator value: **6**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.77**
Disturbance frequency (herb layer): **1.87**
Disturbance severity: **0.76**
Disturbance severity (herb layer): **0.52**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.22**
Grazing pressure: **0.34**
Soil disturbance: **0.3**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae*, QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue, S72 Eastern Mediterranean spiny heath (phrygana)**

Constant species of EUNIS habitats: **S63 Eastern garrigue, S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz