

Bromus japonicus

Habitus and growth type

Plant height [m]: **0.37**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **27.45**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **2.59**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Temperature indicator value: **7.1**

Moisture indicator value: **3.7**

Reaction indicator value: **7.3**

Nutrient indicator value: **4.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.27**

Disturbance frequency (herb layer): **1.98**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.61**
Mowing frequency: **0.59**
Grazing pressure: **0.28**
Soil disturbance: **0.38**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) Sisymbrietea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3D Mediterranean Cupressaceae forest](#)

Constant species of EUNIS habitats: [T3D Mediterranean Cupressaceae forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: 7

Continentality amplitude: 5