

# *Buddleja davidii*

## Habitus and growth type

Plant height [m]: **2.61**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm<sup>2</sup>/mg]: **15.3**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **neophyte**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **6.4**

Moisture indicator value: **5**

Reaction indicator value: **6.6**

Nutrient indicator value: **5**

Salinity indicator value: **0**

## Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.4**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.18**

Grazing pressure: **0.2**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CF \(ROB\) Robinietea](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **5**