

# *Calla palustris*

## Habitus and growth type

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **31.87**

## Fruit, seed and dispersal

Seed mass [mg]: **2.33**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.3**

Temperature indicator value: **5.2**

Moisture indicator value: **9.4**

Reaction indicator value: **6**

Nutrient indicator value: **4.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.58**

Disturbance frequency (herb layer): **1.28**

Disturbance severity: **0.41**

Disturbance severity (herb layer): **0.2**

Mowing frequency: **0.09**

Grazing pressure: **0.11**

Soil disturbance: **0.11**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [IA \(ALN\) \*Alnetea glutinosae\*](#), [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T15 Broadleaved swamp forest on non-acid peat](#)

Constant species of EUNIS habitats: [T15 Broadleaved swamp forest on non-acid peat](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Forest**

## Distribution

Continentality: **7**

Continentality amplitude: **4**