

Camelina rumelica

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **May-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.4**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **6**

Moisture indicator value: **4.2**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2**

Disturbance frequency (herb layer): **2**

Disturbance severity: **0.74**

Disturbance severity (herb layer): **0.74**

Mowing frequency: **0.38**

Grazing pressure: **0.23**

Soil disturbance: **0.55**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**