Campanula cervicaria

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **20.13**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.5**
Temperature indicator value: **5.6**
Reaction indicator value: **7.6**
Nutrient indicator value: **3.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.28**
Disturbance frequency (herb layer): **1.47**
Disturbance severity: **0.6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.22**
Grazing pressure: **0.22**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz