Campanula drabifolia

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **11**
Moisture indicator value: **2**
Reaction indicator value: **8.3**
Nutrient indicator value: **5.5**
Salinity indicator value: **1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***, **KC (ASP) *Asplenietea trichomanis***

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz