

Campanula scheuchzeri

Habitus and growth type

Plant height [m]: **0.23**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **1.72**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.09**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **2.6**

Moisture indicator value: **4.7**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.64**

Disturbance frequency (herb layer): **1.15**

Disturbance severity: **0.4**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.1**
Grazing pressure: **0.24**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) *Calluno-Ulicetea*](#), [CT \(TRI\) *Juncetea trifidi*](#), [CU \(SES\) *Elyno-Seslerietea*](#), [KC \(ASP\) *Asplenetea trichomanis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R23 Mountain hay meadow](#), [R43 Temperate acidophilous alpine grassland](#), [S21 Subarctic and alpine dwarf Salix scrub](#), [S22 Alpine and subalpine ericoid heath](#), [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#), [S26 Subalpine Pinus mugo scrub](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#)

Constant species of EUNIS habitats: [R23 Mountain hay meadow](#), [R41 Snow-bed vegetation](#), [R43 Temperate acidophilous alpine grassland](#), [S21 Subarctic and alpine dwarf Salix scrub](#), [S22 Alpine and subalpine ericoid heath](#), [S24 Subalpine genistoid scrub of the Amphi-Adriatic region](#), [S26 Subalpine Pinus mugo scrub](#), [T34 Temperate subalpine Larix, Pinus cembra and Pinus uncinata forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Heathland, Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **3**