

Campanula trachelocalycina

Habitus and growth type

Plant height [m]: **0.56**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.02**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4**

Temperature indicator value: **5.5**

Moisture indicator value: **4**

Reaction indicator value: **5**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.01**

Disturbance frequency (herb layer): **1.31**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.1**

Mowing frequency: **0**

Grazing pressure: **0.21**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**