

Capsella bursa-pastoris

Habitus and growth type

Plant height [m]: **0.22**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **23.79**

Flower

Flowering period: **January-December**

Fruit, seed and dispersal

Seed mass [mg]: **0.11**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **5.2**

Moisture indicator value: **4.7**

Reaction indicator value: **6.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.79**

Disturbance frequency (herb layer): **2.07**
Disturbance severity: **0.77**
Disturbance severity (herb layer): **0.77**
Mowing frequency: **0.65**
Grazing pressure: **0.21**
Soil disturbance: **0.66**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QA \(PAR\) *Papaveretea rhoeadis*](#), [QB \(SIS\) *Sisymbrietea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V11 Intensive unmixed crops](#), [V12 Mixed crops of market gardens and horticulture](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#), [V15 Bare tilled, fallow or recently abandoned arable land](#), [V35 Trampled mesophilous grassland with annuals](#), [V37 Annual anthropogenic herbaceous vegetation](#)

Constant species of EUNIS habitats: [N1E Black Sea broad-leaved coastal dune forest](#), [V11 Intensive unmixed crops](#), [V12 Mixed crops of market gardens and horticulture](#), [V13 Arable land with unmixed crops grown by low-intensity agricultural methods](#), [V15 Bare tilled, fallow or recently abandoned arable land](#), [V34 Trampled xeric grassland with annuals](#), [V35 Trampled mesophilous grassland with annuals](#), [V37 Annual anthropogenic herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**, **Scrub**, **Forest**, **Sparsely vegetated (incl. rock and scree)**, **Synanthropic**

Distribution

Continentality amplitude: 7