

Cardamine parviflora

Habitus and growth type

Plant height [m]: **0.17**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **34.66**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.1**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **6.3**

Moisture indicator value: **7.4**

Reaction indicator value: **7.2**

Nutrient indicator value: **6.4**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.16**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.48**

Grazing pressure: **0.33**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[OC \(ISO\) *Isoëto-Nanojuncetea*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality: **6**

Continentality amplitude: **5**