

Carduus tmoleus

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **6.6**

Moisture indicator value: **3.9**

Reaction indicator value: **6.5**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.06**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.06**

Grazing pressure: **0.34**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FR \(DAP\) Daphno-Festucetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#), [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [R1K Balkan and Anatolian oromediterranean dry grassland](#), [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub**