

Carex canescens

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **25.6**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.37**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **4.3**

Moisture indicator value: **9.1**

Reaction indicator value: **3**

Nutrient indicator value: **1.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.66**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.37**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.21**

Grazing pressure: **0.14**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [IA \(ALN\) *Alnetea glutinosae*](#), [PA \(SCH\) *Scheuchzeria palustris*-*Caricetea fuscae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q22 Poor fen](#), [Q24 Intermediate fen and soft-water spring mire](#)

Constant species of EUNIS habitats: [Q22 Poor fen](#), [Q24 Intermediate fen and soft-water spring mire](#), [T15 Broadleaved swamp forest on non-acid peat](#), [T16 Broadleaved mire forest on acid peat](#), [T3K Picea mire forest](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Scrub, Forest**

Distribution

Continentality amplitude: **7**