

# *Carex cespitosa*

## Habitus and growth type

Plant height [m]: **0.48**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **18.68**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.62**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.5**

Moisture indicator value: **8.7**

Reaction indicator value: **5.6**

Nutrient indicator value: **4**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.71**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.47**

Grazing pressure: **0.17**

Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#), [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#), [PA \(SCH\) \*Scheuchzerio palustris-Caricetea fuscae\*](#)

### Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

## Distribution

Continentality: **7**

Continentality amplitude: **5**