

Carex cuprina

Habitus and growth type

Plant height [m]: **0.58**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **18.3**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **1.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.4**

Moisture indicator value: **8.1**

Reaction indicator value: **6.7**

Nutrient indicator value: **5.6**

Salinity indicator value: **1.3**

Disturbance indicator values

Disturbance frequency: **1.1**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.44**

Mowing frequency: **0.57**

Grazing pressure: **0.39**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

Constant species of EUNIS habitats: [R34 Submediterranean moist meadow](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Grassland (non-alpine, non-saline),
Saline vegetation, Forest**

Distribution

Continentality amplitude: 7