

# *Carex liparocarpos*

## Habitus and growth type

Plant height [m]: **0.19**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **2.13**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.6**

Temperature indicator value: **6.9**

Moisture indicator value: **2.6**

Reaction indicator value: **7.3**

Nutrient indicator value: **1.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.19**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.41**

Mowing frequency: **0.33**

Grazing pressure: **0.35**

Soil disturbance: **0.21**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FP \(ONO\) Festuco hystricis-Ononidetea striatae](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**