Carex paniculata

Habitus and growth type

Plant height [m]: **0.74**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **16.03**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.77**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**
Temperature indicator value: **5.1**
Moisture indicator value: **9.3**
Nutrient indicator value: **4.4**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.72**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.42**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.34**
Grazing pressure: **0.14**
Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T15 Broadleaved swamp forest on non-acid peat**
Constant species of EUNIS habitats: **T15 Broadleaved swamp forest on non-acid peat**
Dominant species of EUNIS habitats: **Q53 Tall-sedge bed**

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz