

Carex parviflora

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.93**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **1.8**

Nutrient indicator value: **2.6**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.45**

Disturbance frequency (herb layer): **1.52**

Disturbance severity: **0.23**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.02**

Grazing pressure: **0.2**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KE \(THL\) *Thlaspietea rotundifolii*](#), [LA \(HER\) *Salicetea herbaceae*](#), [PB \(OXY\) *Oxycocco-Sphagnetes*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S21 Subarctic and alpine dwarf *Salix* scrub](#)

Broad habitat

Occurrence in broad habitats: **Spring, Mire, Alpine-subalpine and arctic grassland, Heathland, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**

Continentality amplitude: **3**