Carex pseudocyperus

Habitus and growth type

Plant height [m]: **0.57**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **22.91**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.91**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.8**
Temperature indicator value: **5.8**
Moisture indicator value: **9.3**
Reaction indicator value: **6.3**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.68**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.16**
Grazing pressure: **0.11**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OD (PHR) *Phragmito-Magnocaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T15 Broadleaved swamp forest on non-acid peat**
Constant species of EUNIS habitats: **T15 Broadleaved swamp forest on non-acid peat**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Scrub, Forest**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz