Centaurea collina

Habitus and growth type

Plant height [m]: **0.38**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **8.3**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **4**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.44**
Disturbance frequency (herb layer): **2**
Disturbance severity: **0.74**
Disturbance severity (herb layer): **0.72**
Mowing frequency: **0.87**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.2**
Soil disturbance: **0.58**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea*, QF (ART) *Artemisietea vulgaris***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz