

Centaurea nigra

Habitus and growth type

Plant height [m]: **0.54**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.43**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.33**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5**

Moisture indicator value: **4.6**

Reaction indicator value: **4.2**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.69**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **1.02**

Grazing pressure: **0.3**

Soil disturbance: **0.3**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**