

# ***Centaurea nigrescens***

## **Habitus and growth type**

Plant height [m]: **0.66**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Seed mass [mg]: **1.88**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7.8**

Temperature indicator value: **5.7**

Moisture indicator value: **4.8**

Reaction indicator value: **5.8**

Nutrient indicator value: **4.7**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.93**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.59**

Grazing pressure: **0.2**

Soil disturbance: **0.25**

## **Habitat and sociology**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

## **Distribution**

Continentality: **5**

Continentality amplitude: **3**