Centaurea sarfattiana

Habitus and growth type

Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.2**
Temperature indicator value: **6.6**
Moisture indicator value: **3.2**
Reaction indicator value: **5.5**
Nutrient indicator value: **2.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.21**
Disturbance frequency (herb layer): **2.21**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0**
Grazing pressure: **0.4**
Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S74 Central Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **S74 Central Mediterranean mountain hedgehog-heath**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz