

# ***Centaurea subtilis***

## **Habitus and growth type**

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **April-June**

## **Fruit, seed and dispersal**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **2**

Reaction indicator value: **7**

Nutrient indicator value: **1**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **1.25**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.61**

Grazing pressure: **0.4**

Soil disturbance: **0.15**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**