Centaurium tenuiflorum

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **7**
Reaction indicator value: **7**
Nutrient indicator value: **3.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.05**
Disturbance frequency (herb layer): **1.58**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.27**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea*, MF (JUN) *Juncetea maritimi***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal cliff, Wetland, Saline vegetation, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz